

Southern Connecticut State University
Lobster-Aging Proposal
Submitted to the State of Maine

Principal Investigators: Dr. Sean Patrick Grace, Dr. Sarah Crawford, Southern Connecticut State University, Colleen Giannini, Connecticut Department of Environmental Protection

I. Introduction

This research will determine the age of American lobster (*Homarus americanus*) in the Gulf of Maine (GOM) from multiple orbital carapace lengths (mm). The purpose of this research is to document the age and levels of the aging pigment lipofuscin in lobsters from the GOM. It will also aid in the management of this resource. Recent studies on western rock lobster (*Panulirus cygnus*) indicate that the aging technique proposed in this research would allow independent assessment of population parameter estimates and has shown that most juvenile lobsters are 3-5 years old immediately prior to recruitment (Sheehy 1998). Another study on the European lobster (*Homarus gammarus*) yielded reproducible catch age structures with year-class resolution. In addition, a recent study performed on the European Lobster demonstrated that lipofuscin quantification produced more accurate estimates of age in the species than carapace length (Huglem *et. al.* 2005). This finding has important implications for stock assessments that employ traditional models which include age as an input parameter (Sheehy 1996).

II. Goals

The goals of this research are to:

- a) determine the age of American Lobsters from the Gulf of Maine population using lipofuscin accumulation in the olfactory lobe cell mass in the brain.
- b) Compare the age structure of wild caught American Lobster from Long Island sound to that of the age structure determined for wild caught American Lobster from Gulf of Maine. The latter, Long Island Sound data will be provided to Drs. Grace and Crawford from the Connecticut Department of Environmental Protection, Project Director Colleen Giannini.

III. Objectives

1. Establish levels of lipofuscin accumulation in the population of American lobster from the Gulf of Maine.
2. Correlate lipofuscin concentrations from wild Gulf of Maine lobsters with age, using data derived from pigment accumulation analyses from known age lobsters.

IV. Study Design

The approach employed in this study will be to establish lipofuscin levels from wild lobsters of unknown age from the Gulf of Maine using levels previously established for laboratory raised,

known age animals obtained from various hatcheries. Correlation of pigment levels with age will be determined and compared with published literature. This correlation may then serve as the means by which age can be associated with pigment accumulation in the population of wild lobsters from the Gulf of Maine.

V. Procedures/Experimental Design

The project will comprise the two tasks listed below:

Task 1: Fluorescence examination of Gulf of Maine lobsters

The bidder will isolate tissue samples from the esophageal ganglia of Gulf of Maine lobsters. A maximum of 150 lobsters will be purchased from commercial fishermen from the Gulf of Maine by SCSU, specifically, Drs. Sean Grace and Sarah Crawford. The bidder will purchase 10-20 lobsters from each of the following size class ranges (mm carapace length): 82.5-90.4mm, 90.5-98.4mm, 98.5-106.4mm, 106.5-114.4mm, 114.5-122.5mm, 122.5-130.4mm, and >130.5mm. Rationale for the size ranges chosen is an effort to capture the possibility of 5-8 age classes per molt increment based on research performed by Sheehy (1996) in the United Kingdom. Estimation of approximate prices itemized in the budget were determined using the Atlantic States Marine Fisheries Commission American Lobster Stock Assessment 2000, where the length/weight relationships were published (i.e. 127mm (5") lobster = ~ 4.2 lbs). The bidder will purchase, dissect, store and transport Gulf of Maine lobster brains. Laboratory processing and confocal microscopy will take place at SCSU to determine lipofuscin levels in the neural tissue. Up to five sections (5-7 μ m) of the cell mass will be analyzed for each lobster. Each of these sections will quantify the percent of the olfactory lobe cell mass occupied by lipofuscin, the density of lipofuscin granules in the olfactory lobe cell mass, and the mean diameter of the lipofuscin granules.

Task 2: Data Analyses/Written Reports/Work Schedules

The bidder will complete analyses of the pigment data gathered in Task 1 to determine if size and pigment accumulation are related as previously reported in similar studies from published literature. If so, these data will be used to correlate pigment accumulation in wild Gulf of Maine lobsters with known age lobsters whose lipofuscin concentrations have already been established. If no relationship between pigment accumulation and size is found to exist in Task 1, the bidder will provide analyses of concentrations of lipofuscin in wild Gulf of Maine lobsters by orbital carapace length, information that currently does not exist in published literature for this species.

Work Schedules are delineated and provided in the Milestone Chart, Attachment B. Final report of all completed work will be provided on/before May 31, 2007.

VI. CT DEP Responsibilities

Colleen Giannini, Project Director, American Lobster Project, CT DEP Marine Fisheries Division will provide data on lipofuscin concentration of known age American lobster purchased from hatcheries. These samples were fixed and stored in 10% buffered formalin and maintained in labeled air-tight vials until delivered to the SCSU's histology laboratory for processing. All samples were analyzed using the protocol described above and these controls will be used to correlate the results from lobsters collected from the Gulf of Maine. Letter of collaboration from Colleen Giannini is attached. In addition, a letter from support for the proposed research from

Penelope Howell, CT DEP Marine Fisheries Biologist, American Lobster project and chair of the Atlantic States Marine Fisheries Commission Lobster Technical Committee is attached.

VII. Literature Cited

Huglem I., Belchier M., and Svasand T. (2005) 'Age determination of European Lobsters (*Homarus gammarus* L.) by histological quantification of lipofuscin' *J. Crust. Biol.* 25:95-99.

Sheehy M. R. J., Shelton P. M. J., Wickins JF, Belchier M and Gaten E (1996) 'Ageing the European lobster *Homarus gammarus* by the lipofuscin in its eyestalk ganglia' *Mar Ecol Prog Ser* 143:99-111.

Sheehy M, Caputi N, Chubb C and Belchier M. (1998) 'Use of lipofuscin for resolving cohorts of western rock lobster (*Panulirus cygnus*)' *Can J Fish Aquat Sci* 55:925-936.

Simpson D., *et al* (2005) 'Assessment and Monitoring of the American Lobster Resource and Fishery in Long Island Sound Marine Finfish Survey', Job 7 Annual Report, CT DEP Marine Fisheries Division, Old Lyme, CT. 56 pp.

Stock Assessment Report No. 06-03 (Supplement) of the Atlantic States Marine Fisheries Commission (*American Lobster Stock Assessment for peer review*) January 2006.

Wahle RA, Tully O and O'Donovan V (1996) 'Lipofuscin as an indicator of age in crustaceans: analysis of the pigment in the American lobster *Homarus americanus*' *Mar Ecol Prog Ser* 138:117-123

VIII. References Reviewed

Belchier M., Shelton P. M. J. and Chapman C. J. (1994) 'The identification and measurement of fluorescent age pigment abundance in the brain of a crustacean (*Nephrops norvegicus*) by confocal microscopy' *Comp. Biochem. Physiol.*, 108B:157-164.

Blaustein D. N., Derby C. D., Simmons R. B. and Beall A. C. (1988) 'Structure of the brain and medulla terminalis of the spiny lobster *Panulirus argus* and the crayfish *Procambarus clarkii*, with an emphasis on olfactory center' *J. Crustacean Biol.*, 8:493-519.

Hack M. H. and Helmy, F. M. (1983) 'The melanins and lipofuscin' *Comp. Biochem. Physiol.*, 76B:399-407.

Hammer C. and Braum, E. (1988) 'Quantification of age pigments (lipofuscin)' *Comp. Biochem. Physiol.*, 90B:7-17.

Hill K. T. and Womersley, C. (1991) 'Critical aspects of fluorescent age-pigment methodologies: modification for accurate analysis and age assessments in aquatic organisms' *Marn. Biol.*, 109:1-11.

- Hirche H. J. and Anger, K. (1987) 'The accumulation of age pigments during the larval development of the spider crab, *Hyas araneus* (Decapoda, Majidae)' *Comp. Biochem. Physiol.*, 88B:777-782.
- Incze L. S. and Wahle RA (1991) 'Recruitment from pelagic to early benthic phase in lobsters *Homarus americanus*' *Mar Ecol Prog Ser* 79:77-87.
- MacKenzie B. R. (1988) 'Assessment of temperature effects on interrelationships between stage durations, mortality and growth in laboratory-reared *Homarus americanus*. Milne-Edwards larvae' *J. Exp. Mar. Bio. Ecol.*, 116:87-98.
- Nelson K., Hedgecock D., Borgeson W., Johnson E., Daggett R. and Aronstein D (1980) 'Density-dependent growth inhibition in lobsters, *Homarus americanus* (Decapoda, Nephropidae)' *Biol Bull* 159:162-176.
- Nicol S. (1987) 'Some limitations on the use of the lipofuscin ageing technique' *Mar Biol* 93:609-614.
- O'Donovan V. and Tully O. (1996) 'Lipofuscin (age pigment) as an index of crustacean age: correlation with age, temperature and body size in cultured juvenile *Homarus gammarus* L.' *J Exp Mar Bio Ecol.*, 207:1-14.
- Sheehy M. R. J. and Bannister R. C. A. (2002) 'Year-class detection reveals climatic modulation of settlement strength in the European lobster, *Homarus gammarus*.' *Can J Fish Aquat Sci* 59:1132-1143.
- Sheehy M. R. J. and Shelton, P. M. J. (2001) 'Use of lipofuscin age pigment for age determination of lobsters' Marine Fisheries R & D Final Report MF0215/CSA 2596. Ministry of Agriculture Fisheries and Food, London, U. K.
- Sheehy M. R. J., Greenwood, J. G. and Fielder, D. R. (1994) 'More accurate chronological age determination of crustaceans from field situations using the physiological age marker, lipofuscin' *Mar Biol* 121:237-245.
- Sheehy M. R. J. (1992) 'Lipofuscin age pigment accumulation in the brains of aging field and laboratory reared crayfish *Cherax quadricarinatus* (von Martens) (Decapoda: Parastacidae)' *J. Exp. Mar. Biol. Ecol.* 161:79-89.
- Sheehy M. R. J. (1990) 'Individual variation in and the effect of rearing temperature and body size on the concentration of fluorescent morphological lipofuscin in the brains of freshwater crayfish, *Cherax cuspidatus* (Crustacea: Parastacidae)' *Comp. Biochem. Physiol* 96a:281-286.
- Sheehy M. R. J., (1990) 'The widespread occurrence of fluorescent morphological lipofuscin in the crustacean brain' *J. Crust. Biol* 104:613-622.

Sheehy M. R. J., (1990) 'The potential of morphological lipofuscin age pigment as an index of crustacean age' *Mar. Biol* 107:439-442.

Sheehy M. R. J. (1989) 'Crustacean brain lipofuscin: an examination of the morphological pigment in the freshwater crayfish *Cherax cuspidatus*' *J. Crustacean Biol.* 9:387-391.

Sohal R. S. (1987) 'Quantification of lipofuscin: a critique of the current methodology' *Adv. Biosciences* 64:85-91.

Tully O., O'Dovovan V. and Fletcher D. (2000) 'Metabolic rate and lipofuscin accumulation in juvenile European lobster (*Homarus gammarus*) in relation to simulated seasonal changes in temperature' *Mar. Biol* 137:1031-1040.

Van de lest C. H. A., Versteeg E. M. M., Veerkamp J. H. and Vankuppevelt T. H.(1995) 'Elimination of autofluorescence in immunofluorescence microscopy with digital image processing' *J. Histochem. Cytochem* 43:727-730.

Wahle R. A. and Steneck R. S. (1991) 'Recruitment habitats and nursery grounds of the American lobster *Homarus americanus*: a demographic bottleneck?' *Mar. Ecol. Prog. Ser* 69:231-243.

Grant Proposal Specifics: As outlined in the instructions for proposals.

1. The proposal must include detail on the experience and knowledge required as previously stated, including a description of experience.

Description of Southern Connecticut State University's Experience:

Previous awards have been granted to Southern Connecticut State University to support the Master's thesis research of Connecticut Department of Environmental Protection (CT DEP) biologist, Colleen Giannini. These awards have resulted in the purchase and use of a BIORAD Confocal Microscope, Laserssharp 2000 imaging software and related laboratory supplies necessary for histological preparation and paraffin wax sectioning. In addition, the brains of 50 laboratory raised, known age lobsters, ranging in age from 6 months to 5+years, have been analyzed. Lipofuscin concentrations in these animals were quantified using a National Institute of Health (NIH) analysis package and provide the benchmark by which lipofuscin concentrations from wild animals can be correlated with age. SCSU has a sophisticated histological facility dedicated to the ongoing research of fluorescence microscopy and cell imaging. The University recently obtained state of the art digital imaging equipment to compliment their research projects. The investigators have demonstrated specific skills required to isolate the olfactory lobe cell mass from the brain and process the material for lipofuscin detection. Dr. Sean Grace and Dr. Sarah Martinelli are also very familiar with the analytical procedures required to quantify and correlate the laboratory results. They have demonstrated considerable accessibility to the CT DEP's project director, Colleen Giannini, a graduate student under their thesis advisory at SCSU. This association allowed time sensitive samples to be obtained and processed opportunistically, an important factor in meeting the collection requirements to satisfy the contract with the CT DEP. This past history has provided the basis for all future research aimed at lobster aging at SCSU. The equipment is present, the protocol has been established, the experimental design is well tested and the methods are consistent and repeatable.

Bidder's Specific Experience:

Drs. Grace and Crawford have provided the academic advising for the Master's thesis of Colleen Giannini, which involves aging American lobster collected from Long Island Sound. The results of this research have been presented to the Atlantic States Marine Fisheries Commission's American Lobster Stock Assessment, Modeling and Technical Committees in Old Lyme, CT, December 2004, at the Retirement Symposium of Dr. Stann Cobb at the Graduate School of Oceanography at the University of Rhode Island in July 2005, and is scheduled to be presented at the Benthic Ecology Meeting to be held in Quebec, Canada on March 9, 2006. Both Drs. Grace and Crawford have the experience necessary to carry out the histological procedures, perform confocal microscopy and to analyze the digital images of the cell mass. Dr. Grace has also extended the study begun by Colleen Giannini (scheduled for completion in August 2006), by examining the effects of temperature on lipofuscin concentrations in known age animals. The future focus of this research will be the collection of wild lobsters from specific size classes, along a temperature gradient (from Maine to New Jersey) to determine the effects of temperature on natural concentrations of lipofuscin.

Confirmation of award subject to the Freedom of Access Laws of the State of Maine (1 MRSA 401):

We understand that our bid will be available for public viewing following the bid award.

PROJECT PROPOSAL SUMMARY

Project Title: Age Determination of American Lobster in the Gulf of Maine (GOM) using lipofuscin concentrations in the olfactory lobe cell mass of the brain.

Summary of Scope of Work (see above for complete details)

Task 1 – Determine lipofuscin concentrations for wild GOM lobsters

The bidder will quantify lipofuscin accumulations from a minimum of 100 wild caught American lobsters collected from GOM waters by commercial fishermen. A minimum of 100 up to a maximum of 150 lobsters will be purchased from fishermen by Southern Connecticut State University, specifically, Dr. Sean Grace's lab. The bidder will dissect and fix the tissues immediately in 10% neutral buffered formalin, dehydrate, clear and embed the tissues, serially section the brains at 5-7 μ m and slide mount as described in the literature (Simpson *et al* 2005). The bidder will then use confocal microscopy to detect the naturally fluorescent pigment and quantify its presence using digital image analysis (ImageJ™).

Task 2 – Compare lipofuscin concentrations between known age and wild caught.

The bidder will use the results obtained from Task 1 and compare them to the published literature of similar studies done on known age American lobster and also on other crustaceans. This comparison will determine the appropriateness of correlating the results with accumulation levels from wild animals to determine age.

Deliverables

The bidder will complete analyses of the data gathered in Tasks 1 and 2 to determine if the correlation between age and lipofuscin quantities in known age animals can be used to correlate lipofuscin levels in wild GOM lobsters with age.

2. The following individuals are designated to serve as Project Directors:

Southern CT State University

Dr. Sean Patrick Grace
Biology Department
Southern CT State University
501 Crescent Street
New Haven, CT 06515-1355
203-392-6216

Dr. Sarah Crawford
Biology Department
Southern CT State University
501 Crescent Street
New Haven, CT 06515-1355
203-392-6215

3. The following individuals are designated to serve as contacts for business matters:

For the University:

Delinda Conte
Assistant Director of Pre-Award Services
Sponsored Programs and Research (SPAR)
Southern Connecticut State University
501 Crescent Street
New Haven, CT 06515-1355
1-203-392-5066

4. Total project costs for the period of the Agreement shall not exceed \$7,049.00.

SOUTHERN CT STATE UNIVERSITY

by _____
Authorized signature

Typed Name and Title

Date

ATTACHMENT A
Cost Proposal

1. Provide a fixed cost proposal to deliver the research and final product previously detailed.

Proposed Costs: Bidders would use monies awarded for the following items, please note that no salary is being requested. Drs. Grace and Crawford have access to excellent graduate students, whom will assist in the dissection, laboratory processing and digital imaging and analysis of lobster tissues.

Item 1: Laboratory costs:

Histological processing equipment.	\$2,000.00
Laboratory chemicals and sectioning supplies.	\$1,500.00

Item 2: Travel Expenses (to Maine to collect, dissect and store neural tissue.) . . . \$600.00

Item 3: GOM Lobster purchases (estimating wholesale cost at \$6.00/lb):

10-20 lobsters 82.5-90.4mm CL (approximately 2.7lbs) x \$6.00/lb	\$326.00
10-20 lobsters 90.5-98.4mm CL (approximately 2.9lbs) x \$6.00/lb.	\$358.00
10-20 lobsters 98.5-106.5mm CL (approximately 3.2lbs) x \$6.00/lb	\$390.00
10-20 lobsters 106.5-114.4mm CL (approximately 3.8lbs) x \$6.00/lb	\$422.00
10-20 lobsters 114.5-122.4mm CL (approximately 4.0lbs) x \$6.00/lb	\$453.00
10-20 lobsters 122.5-130.4mm CL (approximately 4.2lbs) x \$6.00/lb	\$485.00
10-20 lobsters >130.5mm CL (approximately 4.3lbs) x \$6.00/lb	\$515.00

Total \$7,049.00